



GETTING STARTED WITH AFREZZA®

THE ONLY ULTRA-RAPID ACTING INHALED INSULIN FOR ADULTS LIVING WITH TYPE 1 OR TYPE 2 DIABETES



Indications and Limitations of Use

Afrezza® (insulin human) Inhalation Powder is a man-made rapid acting insulin breathed in through your lungs (inhaled) and is used to control high blood sugar in adult patients with diabetes mellitus. AFREZZA is available by prescription only.

- AFREZZA is not for use in place of long-acting insulin. AFREZZA must be used with long-acting insulin in people who have type 1 diabetes.
- AFREZZA is not for use to treat diabetic ketoacidosis.
- It is not known if AFREZZA is safe and effective for use in people who smoke. AFREZZA is not for use in people who smoke or have recently stopped smoking (less than 6 months).
- It is not known if AFREZZA is safe and effective in children under 18 years of age.

AFREZZA can cause serious side effects including sudden lung problems (bronchospasms). Do not use AFREZZA if you have long-term (chronic) lung problems such as asthma or chronic obstructive pulmonary disease (COPD). Before starting AFREZZA, your healthcare provider will give you a breathing test to check how your lungs are working.

Please see Important Safety Information for Afrezza on page 12 or on www.afrezza.com. The complete Medication
Guide and Instructions for Use can be found on www.afrezza.com.



TAKING CONTROL WITH AFREZZA®

As someone living with type 1 or type 2 diabetes who depends on mealtime insulin, you know how important flexibility can be. Afrezza is the only ultra rapid-acting inhaled insulin that works on your schedule to deliver mealtime management in the moment.



NO PLANNING

Afrezza lets you inhale your insulin right when food arrives—even unexpectedly—so you can be spontaneous but still in control.



NO WAITING

Afrezza lowers blood sugars rapidly, similar to the insulin response of a person without diabetes.



NO NEEDLESTICKS

Afrezza is an innovative solution that gives you mealtime glucose control without the need for mealtime injections.





WHAT IS AFREZZA®?

MEALTIME ON YOUR TIME

- Inhaled insulin that gets into your bloodstream in less than 1 minute
- Starts controlling blood sugar in about 12 minutes

CONTROL YOU CAN COUNT ON

- Eliminates over 1,000 mealtime injections per year*
- Proven mealtime control in studies of more than
 3,000 people living with type 1 or type 2 diabetes

Since Afrezza leaves the body within 1.5 to 3 hours,[†] depending on the dose, it gives you the option to take additional doses at 1 and 2 hours after meals, if needed.

^{*}Assumes 1 needle per meal, 3 meals per day for 365 days, rounded down.

⁺For the 4-unit and 12-unit cartridges, respectively.





FILLING YOUR PRESCRIPTION

Here's a step-by-step guide to filling your first Afrezza® prescription:

- Before you visit the pharmacy, be sure to download your Afrezza Savings Card at www.AfrezzaSavingsCard.com
- 2. Take both your prescription and savings card to the pharmacy, and be prepared to give them at least 48 hours to fill your order as some locations may need to have Afrezza shipped to them

CONTROL YOUR COSTS WITH THE AFREZZA SAVINGS CARD*



WE ARE HERE TO HELP

Contact your MannKind Support Specialist with any issues related to your Afrezza prescription.

We can help you explore options to get support if your health plan or pharmacist tells you Afrezza is not covered on your current plan.

The MannKind Customer Experience Center: 1-833-623-4843

*Eligibility criteria and maximum benefit limitations apply.
Please see www.AfrezzaSavingsCard.com for complete details.

TAKING AFREZZA®



With these simple steps, learn how to manage your mealtime glucose levels.



EFFECTIVE DOSING

Like all insulin therapy, monitor the effect Afrezza has on you, and reach out to your doctor if you feel like you need to change your dosing.



TYPICAL DOSING AT MEALTIME

In clinical trials, most patients received 12-24 Afrezza insulin units per meal.



CONVERSION CONSIDERATIONS

Based on clinical trials, consider a ~1.5x conversion from your injected dose of insulin to your inhaled dose of Afrezza to achieve a comparable glucose effect. You may need to adjust your dose, up or down, based on your blood sugar levels and how Afrezza works for you.

Frequently check your blood glucose levels to see how Afrezza is working for you.

TAKING AFREZZA®



With these simple steps, learn how to manage your mealtime glucose levels.

DOSING WITH CARTRIDGES

If you've had experience with mealtime injections, use this starting dose conversion chart to help determine the right amount of Afrezza insulin units you need.

INJECTED	AFREZZA	# OF CARTRIDGES NEEDED		NEEDED
MEALTIME INSULIN DOSE	STARTING DOSE	4 UNIT BLUE	8 UNIT GREEN	12 UNIT YELLOW
UP TO 4	4 UNITS			
5-8 units	8 UNITS			
9-12 units	12 UNITS		+	OR D
13-16 units	16 UNITS			>
17-20 units	20 UNITS			+
21-24 UNITS	24 UNITS			

Consider a 1.5 conversion from injectable insulin to Afrezza units to achieve a comparable effect.

TAKING AFREZZA®



With these simple steps, learn how to manage your mealtime glucose levels.

MAKING ADJUSTMENTS TO MAXIMIZE RESULTS

As you begin your new therapy, be sure to test your blood glucose levels frequently, to track how Afrezza is working for you. Discuss your results with your healthcare provider so you can work together to adjust the Afrezza dose to what works for you.

AFREZZA DOSING PER MEAL

Work with your doctor to determine the dosing regimen that's right for you, then record it here:

STEP 1: Take prescribed Afrezza dose

BREAKFAST	LUNCH	DINNER	SNACK
units	units	units	units

STEP 2: Check blood sugar to see how Afrezza is working for you

_ hour(s) after beginning the meal*

STEP 3: Follow-up dose, if needed

AFTER-MEAL BLOOD SUGAR	FOLLOW-UP DOSE
Target blood sugar: Between mg/dL and mg/dL	No follow-up dose needed
Greater than mg/dL	Take units

STEP 4: Adjust dose, if needed

- If your after-meal blood sugar remains above your target blood sugar range for _______ days consult your doctor to consider increasing your dose in Step 1
- If your after-meal blood sugar is lower than your target range, consult your doctor
- Continue adjusting until your after-meal blood sugar is consistently within your target

From changes in diet and exercise to stress levels and medications, life is constantly changing. Fortunately, Afrezza can be adapted to change with it, with a variety of cartridge sizes (4-unit, 8-unit, and 12-unit cartridges) to help manage any necessary adjustments.

^{*}American Diabetes Association suggests 1–2 hours after beginning of the meal. The American Association of Clinical Endocrinologists suggests 2 hours after beginning of the meal.

HOW TO USE AFREZZA®



A STEP-BY-STEP GUIDE TO YOUR FIRST DOSE

Before you take your first dose, make sure to learn proper inhalation techniques and how Afrezza works. You can discuss instructions and receive a virtual training demonstration with a MannKind Patient Support Guide.

Call 1-833-623-4843 to learn more. Watch the How to Use Afrezza video at www.Afrezza.com.

STEP 1: SELECT THE CARTRIDGE FOR YOUR DOSE



4 UNITS:

If your Afrezza dose is 4 units, use 1 blue cartridge.



8 UNITS:

If your Afrezza dose is 8 units, use 1 green cartridge.




12 UNITS:


If your Afrezza dose is 12 units, use 1 yellow cartridge.


If your prescribed Afrezza dose is more than 12 units, you will need to use more than 1 cartridge to get your right dose.


STEP 2: LOAD A CARTRIDGE

- HOLD INHALER: Hold the inhaler level in one hand with the white mouthpiece on the top and purple base on the bottom.
- **OPEN INHALER:** Open the inhaler by lifting the white mouthpiece to a vertical position.
- PLACE CARTRIDGE: Hold the cartridge with the cup facing down. The pointed end of the cartridge should line up with the pointed end in the inhaler. Place cartridge into the inhaler, making sure it lies flat in inhaler.
- **KEEP LEVEL:** Now that the cartridge is loaded, keep the inhaler level from this point forward to avoid loss of drug powder. Do not turn it upside down, shake, or drop, as this could cause a loss of drug powder.
- CLOSE INHALER: Lower the mouthpiece to close the inhaler (this will open the drug cartridge). You should feel a snap when the inhaler is closed.











HOW TO USE AFREZZA®




STEP 3: INHALE INSULIN

- **REMOVE COVER:** Remove the mouthpiece cover. Keep the inhaler level.
- **EXHALE:** Hold the inhaler away from your mouth and exhale fully.
- **POSITION INHALER:** Keeping your head and inhaler level, place the mouthpiece in your mouth. Close your lips around the mouthpiece to form a seal. Tilt the inhaler downward while keeping your head level as shown.
- INHALE DEEPLY AND HOLD BREATH: With your mouth closed around the mouthpiece, inhale deeply through the inhaler. Hold your breath for as long as comfortable and at the same time remove the inhaler from your mouth. Exhale and breathe normally.








STEP 4: REMOVE CARTRIDGE AND DISCARD

- **REPLACE COVER:** Put the mouthpiece cover back onto the inhaler.
- OPEN INHALER: Open the inhaler by lifting the white mouthpiece.
- **REMOVE CARTRIDGE:** Remove the cartridge from the purple base.
- **DISCARD:** Throw away the cartridge in your regular household trash.

Repeat steps 2 through 4 for each Afrezza cartridge you need for your dose.











STORAGE AND HANDLING

HOW TO TAKE CARE OF YOUR AFREZZA® INHALER AND INSULIN CARTRIDGES

IN USE: **ROOM TEMPERATURE STORAGE**

Reference the chart below for instructions on taking care of your inhaler and opened foil packages.

OPENED AFREZZA INHALERS	Room Temperature
	Use for up to 15 days from the date of first use. After 15 days, inhaler must be discarded and replaced.
SEALED BLISTER CARDS + STRIPS	Room Temperature
	Must be used within 10 days
OPENED STRIPS	Room Temperature
	Must be used within 3 days

Do not put a blister card or strip back into the refrigerator after being stored at room temperature.

NOT IN USE: REFRIGERATED STORAGE

Store unopened drug in a refrigerator 36°F-46°F (2°C-8°C).

SEALED FOIL PACKAGES	Refrigerated
afreza	May be used until the expiration date*
SEALED BLISTER CARDS + STRIPS	Refrigerated
	Must be used within 1 month*

*If a foil package, blister card, or strip is not refrigerated, the contents must be used within 10 days.

BEFORE USING YOUR AFREZZA INHALER



Before use, cartridges and inhaler should be at room temperature for 10 minutes.





10 mins.



FREQUENTLY (insulin human) Inhalation Powder ASKED QUESTIONS

Q: WHEN SHOULD I TAKE MY AFREZZA®?

A: Afrezza should be inhaled at the beginning of a meal.

Q: IS IT NORMAL TO SEE POWDER LEFT OVER IN MY INHALER AFTER I'VE INHALED MY AFREZZA DOSE?

A: Yes, a small amount of powder residue in the mouthpiece or cartridge is normal. This does not affect your dose. You can wipe the mouthpiece with a clean, dry cloth to remove moisture or lipstick. Do not wash the Afrezza inhaler.

Q: WHAT SHOULD I DO IF I EXPERIENCE A COUGH?

A: Since Afrezza is an inhaled powder, it is not uncommon to experience cough at the beginning of your treatment. In our clinical studies, cough was mild and typically declined with continued use after 1–4 weeks. As a suggestion, taking Afrezza at room temperature and having a sip of water before and after inhalation may help. If you develop a persistent or recurring cough, or have breathing difficulties while using Afrezza, contact your healthcare provider.

ADDITIONAL QUESTIONS?

Visit **www.Afrezza.com** or contact the MannKind Customer Experience Center at **1-833-623-4843**.

THE MANNKIND CUSTOMER EXPERIENCE CENTER



HERE TO HELP EVERY STEP OF THE WAY

YOUR FIRST PRESCRIPTION

Be sure to **download** the Afrezza® Savings Card **before you visit your pharmacy**, and give them at least 48 hours to fill your prescription as some locations may need to have the product delivered.*

YOUR FIRST DOSE

Remember, Afrezza is different from injectable insulin, delivering ultra-rapid control that may reduce glucose levels faster than you may be accustomed to.

YOUR FIRST FOLLOW-UP APPOINTMENT

Once you and your doctor have decided Afrezza is right for you, be sure to schedule a 2-week follow-up to discuss initial results, and address any issues or adverse reactions.

YOUR CONTINUED SUCCESS

Your doctor will be your best resource as you optimize dosing and work through any common side effects, including coughing, which is generally mild and tends to decline with continued use.

CONTACT US:





EMAIL guida@mannkindcorn.com

WEBSITE www.Afrezza.com

THE MANNKIND CUSTOMER EXPERIENCE CENTER

can help you navigate insurance coverage and explain how to use your Afrezza Savings Card.

THE MANNKIND CUSTOMER EXPERIENCE CENTER

can help you understand what to expect during the first few weeks of treatment and provide instructions on proper inhalation technique.

THE MANNKIND CUSTOMER EXPERIENCE CENTER

can help you develop questions for your healthcare team to make sure you get the most out of your Afrezza experience.

THE MANNKIND CUSTOMER EXPERIENCE CENTER

provides training and support as you need, and helps you identify additional financial programs you may be eligible for that can keep your treatment affordable.

Indications and Limitations of Use

Afrezza® (insulin human) inhalation powder is a man-made rapid acting insulin breathed in through your lungs (inhaled) and is used to control high blood sugar in adult patients with diabetes mellitus. AFREZZA is available by prescription only.

- AFREZZA is not for use in place of long-acting insulin.
 AFREZZA must be used with long-acting insulin in people who have type 1 diabetes.
- AFREZZA is not for use to treat diabetic ketoacidosis.
- It is not known if AFREZZA is safe and effective for use in people who smoke. AFREZZA is not for use in people who smoke or have recently stopped smoking (less than 6 months).
- It is not known if AFREZZA is safe and effective in children under 18 years of age.

Important Safety Information

AFREZZA can cause serious side effects including:

 Sudden lung problems (bronchospasms). Do not use AFREZZA if you have long-term (chronic) lung problems such as asthma or chronic obstructive pulmonary disease (COPD). Before starting AFREZZA, your healthcare provider will give you a breathing test to check how your lungs are working.

Tell your healthcare provider if you:

- Have lung problems such as asthma or COPD
- Have or have had lung cancer
- Are using any inhaled medications
- Smoke or have recently stopped smoking
- Have kidney or liver problems
- Are pregnant, planning to become pregnant, or are breastfeeding.
 AFREZZA may harm your unborn or breastfeeding baby.
- Are taking any other medications, including prescription and over-the-counter medicines, vitamins, or herbal supplements

Before you start using AFREZZA, talk to your healthcare provider about low blood sugar and how to manage it.

The most common side effects of AFREZZA are low blood sugar (hypoglycemia), cough, and sore throat. These are not all the possible side effects of AFREZZA. Call your healthcare provider for medical advice about side effects.

You may report side effects to the FDA at 1-800-FDA-1088 (1-800-332-1088) or www.fda.gov/medwatch.

Please see complete <u>Medication Guide</u> and <u>Instructions</u> <u>for Use</u> on <u>www.afrezza.com</u>

References:

- 1. Afrezza (insulin human) Inhalation Powder Prescribing Information. MannKind Corporation.
- 2. Bode BW, McGill JB, Lorber DL, et al. Inhaled Technosphere insulin compared with injected prandial insulin in type 1 diabetes: a randomized 24-week trial. *Diabetes Care*. 2015;38(12):2266–2273.
- 3. Rosenstock J, Lorber DL, Gnudi L, et al. Prandial inhaled insulin plus basal insulin glargine versus twice daily biaspart insulin for type 2 diabetes: a multicentre randomised trial. *Lancet*. 2010;375(9733):2244–2253.
- 4. Rosenstock J, Franco D, Korpachev V, et al. Inhaled Technosphere insulin versus inhaled Technosphere placebo in insulin-naive subjects with type 2 diabetes inadequately controlled on oral antidiabetes agent. *Diabetes Care*. 2015;38(12):2274–2281.
- 5. Akturk HK, Snell-Bergeon JK, Rewers A, et al. Improved postprandial glucose with inhaled Technosphere insulin compared with insulin aspart in patients with type 1 diabetes on multiple daily injections: the STAT study. *Diabetes Technol Ther.* 2018;20(10):639–647.
- 6. Data on file. MannKind Corporation.

